

[IJPTHer] Editor Decision

1 pesan

Mustofa via Jurnal Ilmiah Universitas Gadjah Mada <noreply-ojs3@ugm.ac.id>

19 Mei 2022 pukul 11.28

Balas Ke: Mustofa <mustofak@ugm.ac.id>

Kepada: Ni Luh Gede Sudaryati <sudaryati@unhi.ac.id>, Ni Made Riris Divayani Aristyantari <ririsdivayani17@gmail.com>, Ni Ketut Ayu Juliasih <juliasih@unhi.ac.id>, I Nyoman Arsana <arsanacita@gmail.com>

Ni Luh Gede Sudaryati, Ni Made Riris Divayani Aristyantari, Ni Ketut Ayu Juliasih, I Nyoman Arsana :

We have reached a decision regarding your submission to Indonesian Journal of Pharmacology and Therapy, "Antibacterial activities of *Averrhoa bilimbi* L. (Oxalidaceae) fruit extract inhibiting *Salmonella typhi*".

Our decision is: Revisions Required

Mustofa
Universitas Gadjah Mada
mustofak@ugm.ac.id

Reviewer A:
Recommendation: Revisions Required

General comment

The manuscript presents the result of the antimicrobial assay of herbal. There are several correction needs to be made to improve the manuscript:

1. Provide the name of the herbal in Bahasa Indonesia or the local name.
2. Provide the information regarding the groups of treatment in the methods section.
3. Table 1 should be completed with the information regarding the group name and the abbreviation mean.
4. figure 1 : please provide the information which disk is P1, p2, p3, K-, K+, etc and write what is the thing that the author want to show by showing the picture

Introduction

The manuscript presents the result of the antimicrobial assay of herbal. There are several correction needs to be made to improve the manuscript:

1. Provide the name of the herbal in Bahasa Indonesia or the local name.
2. Provide the information regarding the groups of treatment in the methods section.
3. Table 1 should be completed with the information regarding the group name and the abbreviation mean.
4. figure 1 : please provide the information which disk is P1, p2, p3, K-, K+, etc and write what is the thing that the author want to show by showing the picture

Methodology

The manuscript presents the result of the antimicrobial assay of herbal. There are several correction needs to be made to improve the manuscript:

1. Provide the name of the herbal in Bahasa Indonesia or the local name.
2. Provide the information regarding the groups of treatment in the methods section.
3. Table 1 should be completed with the information regarding the group name and the abbreviation mean.
4. figure 1 : please provide the information which disk is P1, p2, p3, K-, K+, etc and write what is the thing that the author want to show by showing the picture

Results

The manuscript presents the result of the antimicrobial assay of herbal. There are several correction needs to be made to improve the manuscript:

1. Provide the name of the herbal in Bahasa Indonesia or the local name.
2. Provide the information regarding the groups of treatment in the methods section.
3. Table 1 should be completed with the information regarding the group name and the abbreviation mean.
4. figure 1 : please provide the information which disk is P1, p2, p3, K-, K+, etc and write what is the thing that the author want to show by showing the picture

Discussion

-

Conclusion

-

References

ok

Recomendation

- Accepted, requires minor revision

Reasons for rejected (*Please be specific*)

Additional comments

Reviewer C:

Recommendation: Accept Submission

General comment

Nich to read arti lebih. Scientifically sounds.

Introduction

Good

Methodology

Appropriate.

Results

Accepted

Discussion

Good

Conclusion

Appropriate

References

Appropriate

Recomendation

- Accepted, requires minor revision

Reasons for rejected (*Please be specific*)

Additional comments

Would be best if the author(s) conduct a phytoscreening analysis of the compounds within the sample.

Antibacterial activities of *Averrhoa bilimbi* L. (Oxalidaceae) fruit extract inhibiting *Salmonella typhi*

ABSTRACT IN ENGLISH

Averrhoa bilimbi L. has been used in traditional medicine (*Usada*) to treat typhoid fever. The fruit's flavonoids and triterpenoids are considered to have antibacterial properties. This study is aimed to investigate antibacterial activities of *A. bilimbi* L. (Oxalidaceae) fruit extract in order to inhibit the *Salmonella typhi*. Four groups were treated with *A. bilimbi* L. 25%, 50%, 75%, and 100%. The ciprofloxacin employed as the positive control, while sterile distilled water as the negative control. The extract's inhibitory activity was determined using *S. typhi* bacteria cultured on *Mueller Hinton Agar* (MHA). Kruskal Wallis and U-Mann Whitney tests with a 95 % level of evidence ($p < 0.05$) were employed to analyze the data. The results indicated that *A. bilimbi* L. fruit extract inhibited *S. typhi* following the concentrations as 25% (0 ± 0 mm), 50% ($13,000 \pm 1,414$ mm) strong criterion, 75% ($18,750 \pm 1,500$) strong criterion, and 100% ($20,250 \pm 1,707$ mm) extremely strong criterion. Statistical analysis revealed a statistically significant difference in concentrations of *A. bilimbi* L. fruit extract and *S. typhi* in all groups, with p -value of 0.000 ($p < 0.05$). Thus, the fruit extract possesses antibacterial properties and has the potential to be developed in medication as herbal medicine (*Usada*).

ABSTRACT IN BAHASA

Pengobatan penyakit demam tifoid dapat memanfaatkan bagian tumbuhan sebagai obat tradisional (*usada*) salah satunya buah *Averrhoa bilimbi* L. Kandungan flavonoid dan triterpenoid pada buah diyakini bermanfaat sebagai antibakteri. Penelitian ini berupaya untuk menguji aktivitas antibakteri ekstrak buah *A. bilimbi* L. (Oxalidaceae) dalam menghambat *S. typhi*. Penelitian eksperimental laboratorium dengan rancangan *Post-test Only Control Group Design*. Terdapat 6 kelompok perlakuan ekstrak *A. bilimbi* L. dengan konsentrasi 25%, 50%, 75%, 100%, kontrol positif (*Ciprofloxacin*) dan kontrol negatif (aquadest steril). Daya hambat ekstrak diujikan pada bakteri *S. typhi* yang ditumbuhkan pada media *Mueller Hinton Agar* (MHA). Analisis data menggunakan uji *Kruskal Wallis* dan *U-Mann Whitney* dengan level of evidence 95% ($p < 0,05$). Hasil penelitian diperoleh ekstrak buah *A. bilimbi* L. dalam menghambat *S. typhi* sebesar 25% (0 ± 0 mm), 50% ($13,000 \pm 1,414$ mm) kriteria kuat, 75% ($18,750 \pm 1,500$) kriteria kuat dan 100% ($20,250 \pm 1,707$ mm) kriteria sangat kuat. Hasil uji statistik menunjukkan terdapat perbedaan signifikan antara konsentrasi pemberian ekstrak buah *A.*

bilimbi L. terhadap *S. typhi* di semua kelompok dengan *p-value* 0,000 ($p < 0,05$). Sehingga, ekstrak buah memiliki aktivitas antibakteri dan berpotensi dikembangkan sebagai obat herbal (*Usada*).

Keywords: antibacterial, *Averrhoa bilimbi* L., typhoid fever, *usada*.

INTRODUCTION

Typhoid fever is one of Indonesia's endemic diseases. This disease is caused by feces and urine contamination of water and food, which introduce *Salmonella typhi* bacteria into the human body ¹. The *S. typhi* bacteria assault the digestive and excretory systems, resulting in systemic infections shown by recurrent fever, bacteremia, and inflammation of the intestines and liver ². *Typhoid* fever frequency and incidence have grown in Indonesia, particularly in places with inadequate sanitation, with 21 million cases documented and over 700 deaths ³.

Efforts to manage and control typhoid fever in Indonesia have been made. However, the disease continues to spread in several places. *Typhoid* fever is treated with first-generation antibiotics (*chloramphenicol*, *co-trimoxazole*, *Ciprofloxacin*, and *amoxicillin*) that inhibit *S. typhi* growth. However, long-term use results in antibiotic resistance in the form of *Multi-Drug Resistance Salmonella typhi* (MDR-ST) ^{4,5}. Thus, deliberate measures are required to prevent the emergence of antibiotic resistance by

using plant components with antibacterial properties that are safe, affordable, and capable of inhibiting the development of typhoid fever pathogens ⁶.

Averrhoa bilimbi L. (Oxalidaceae) is a plant species widely employed in Indonesian traditional medicine, particularly in the province of Bali. In Bali, traditional medicine (*Usada*) is derived from *Lontar Usada Taru Pramana* plant. According to the *papyrus*, *Averrhoa bilimbi* L. is an effective medication, particularly for fever relief. So it is critical to perform empirical investigations on the discovery and development of traditional medicine in Bali. According to Kimbal ⁷, Sunti acid content produced from *Averrhoa bilimbi* L. extract fermentation can inhibit *Escherichia coli* Multidrug-Resistant (ECMR). Furthermore, the *ethanolic* extract of *A. bilimbi* L. has antibacterial activity on all MDR (*Multidrug-Resistant*) bacteria with an inhibitory diameter of 10-14.5 mm on agar diffusion wells. It has potential as an antibacterial agent, especially for MDR strains ⁶.

The leaves of *A. bilimbi* L. contain an ethanol compound with an ethanol concentration of 82.82 mg GAE/g DW in the DPPH test and are capable of inhibiting the growth of *Salmonella sp.*, *Escherichia coli*, and *Staphylococcus aureus* bacteria, suggesting that it may be used to prevent

diseases caused by oxidative stress and bacterial infection ⁸. Additionally, the fruit of *A. bilimbi* L. can be used as an antipyretic, lowering fever effectively ⁹. After reviewing the advantages of *Averrhoa bilimbi* L. as traditional medicine and its potential as an antibacterial agent, This study investigates the antibacterial activities of *Averrhoa bilimbi* L. (Oxalidaceae) fruit extract inhibiting *Salmonella typhi*. The results of this study are intended to serve as a scientific reference for the advancement of traditional medicine (*Usada*), particularly for the investigation of the content and inhibitory potential of *A. bilimbi* L. fruit extract against the pathogen that causes *typhoid* fever.

MATERIALS AND METHODS

Study Design

This research used an experimental laboratory design with a *Post-test Only Control Group*. This study included six treatment groups: negative control (K-) (sterile distilled water), positive control (K+) (*Ciprofloxacin*), and *Averrhoa bilimbi* L. concentrations of 25% (P1), 50% (P2), 75% (P3), and 100% (P4), with four replications determined using the *Federer* equation ¹⁰. For six months, the research was carried out at the

Biopesticide Laboratory and Clinical Microbiology Laboratory, Faculty of Medicine, Udayana University.

Approval Study

The research series has been approved by the Department/KSM Clinical Microbiology, Faculty of Medicine, Udayana University or Central General Hospital (RSUP) Sanglah Denpasar with Number 155/UN.14.2.2.VIII.6/2018.

Instruments and materials

The instruments and materials in this study were: 96% ethanol, sterile distilled water, *Ciprofloxacin* antibiotic disk, *Mueller Hinton Agar* (MHA), *SS Agar* media, blank disk, 0.9% NaCl, sterile cotton sticks, water bath, blender, rotary evaporator, glass beaker, petri dish, spirit lamp, incubator, caliper, autoclave, filter paper.

Study procedure

***Mueller Hinton Agar* (MHA) media creation**

Suspension of 38 g of MHA powder in 1 L of distilled water in an Erlenmeyer. Stir until the MHA powder is dissolved. After dissolving, the pH of the media was measured using a pH stick (optimal pH 7.3 ± 0.1). Sterilized in an autoclave at 121°C for 15 minutes. The cold liquid MHA was poured into a petri dish aseptically and then stored.

***Averrhoa bilimbi* L. extract creation**

The *ethanolic* extract of *A. bilimbi* L. fruit was obtained by maceration of *simplicia* powder with 96% ethanol. One hundred grams of *simplicia* powder of *A. bilimbi* L. fruit was soaked (maceration) in 60 mL of 96% ethanol solution for three days while stirring once a day. The results obtained were filtered and re-maceration with 40 mL of 96% ethanol solution and allowed to stand for two days. All the macerated were then concentrated using a rotary evaporator until a thick extract was obtained.

Dilution of a thick extract and preparation of a concentration

The viscous extract of the macerated *A. bilimbi* L. fruit was an antibacterial extract with a concentration of 100%. Each dilution was made as 5 mL and prepared in four glass beakers. Each coded A, B, C, D. Each glass beaker was given a letter code to accommodate: Code A was used to accommodate extracts with a concentration of 100%, and code B was used to accommodate extracts with a concentration of 100%. With a concentration of 75%, code C is used to accommodate extracts with a concentration of 50%, and code D is used to accommodate extracts with 25%. Then each glass beaker was inserted: Code A: 100% extract. Code B: 3.75 mL of *A. bilimbi* L. fruit extract + 1.25 mL of sterile distilled water.

Code C: 2.5 mL of *A. bilimbi* L. fruit extract + 2.5 mL of sterile distilled water. Code D: 1.25 mL of *A. bilimbi* L. fruit extract + 3.75 mL of sterile distilled water. Each dilution was homogenized, and the extract was ready for use.

***Salmonella typhi* bacterial suspension preparation**

One to three oses of *S. typhi* colonies from pure cultures were taken and suspended into a tube containing 3 mL of 0.9% physiological NaCl solution. This suspension is compared to standard *McFarland* turbidity.

Assay of antibacterial activity

S. typhi bacterial suspension was inoculated using a dispersion technique using a sterile cotton swab on *Mueller Hinton Agar* (MHA) media. Discs impregnated with fruit extract of *A. bilimbi* L. with different concentrations were placed on each test medium. *Ciprofloxacin* antibiotic disks were also placed on each test medium as a positive control and sterile distilled water as a negative control. The media was incubated at 37°C for 1×24 h. The zone of inhibition formed around the disc, measured in mm.

Data Analysis

The zone of inhibition is determined by the strength with which the inhibition is carried out, which is expressed in millimeters (mm). Among the criteria for an inhibition zone are the following: 5 mm (weak); 5-10

mm (medium); 10-20 mm (strong); and > 20 mm (very strong). The data from the inhibition zone for *S. typhi* growth were evaluated statistically using the Statistical Package for the Social Sciences (SPSS) Inc. version 21.0 software. The *Kruskal Wallis* test was used to examine the data, followed by the *U-Mann Whitney* test at a 95% confidence level ($p < 0.05$). Tables, figures, and narratives are used to convey all data.

RESULT

Testing the antibacterial activity of *A. bilimbi* L. (Oxalidaceae) fruit extracts at various concentrations in inhibiting *S. typhi* revealed that a concentration of 100% (P4) has a potent inhibitory activity. In comparison, concentrations of 75% (P3) and 50% (P2) have a moderate inhibitory activity, and a concentration of 25% (P1) has no inhibitory activity and is equivalent to negative control (sterile distilled water). The test results indicate that there are data that do not pass the basic assumption test, necessitating a non-parametric test to determine comparability. The *Chi-Square* score is 22.201, corresponding to a probability of 0.05 ($p = 0.000$). Thus, there is a substantial variation in the ability of *A. bilimbi* L. (Oxalidaceae) fruit extract to inhibit *S. typhi* inhibition across all treatment groups. The *U-Mann Whitney* test revealed significant

differences in the majority of treatment groups except for the negative control group with a concentration of 25% (P1) and a concentration of 75% (P3) with a probability value of 100% (P4) ($p>0,05$). The antibacterial activity of *A. bilimbi* L. fruit extract against *S. typhi* is demonstrated in Table 1 and Figure 1.

TABLE 1. Antibacterial activity of *A. bilimbi* L. fruit extract in inhibiting *S. typhi*

Group	Replication				Inhibition Average Zone	Inhibition Criteria	Shapiro Wilk test	Levene Test	Chi-Square	p-value
	I	II	III	IV						
K(+)	33	37	34	33	34,250 ± 1,892a	Very strong	0,086			
K(-)	0	0	0	0	0 ± 0b	-	0			
P1	0	0	0	0	0 ± 0b	-	0			
P2	15	12	12	13	13,000 ± 1,414c	Strong	0,161	0,019	22,201	0,000*
P3	20	17	18	20	18,750 ± 1,500d	Strong	0,224			
P4	21	18	20	22	20,250 ± 1,707d	Very strong	0,850			

Information: * = there is a significant difference ($p<0.05$); Different letter = there is a significant difference ($p<0.05$); Same letter = no significant difference ($p>0.05$); K(-) = Kontrol negatif; K(+) = Kontrol positif; P1 = *A. bilimbi* L. concentrations of 25%; P2 = *A. bilimbi* L. concentrations of 50%; P3 = *A. bilimbi* L. concentrations of 75%; and P4 = *A. bilimbi* L. concentrations of 100%.




FIGURE 1. *S. typhi* bacteria have been inoculated on *Mueller Hinton Agar* (MHA) media with the addition of discs according to the concentration of the treatment. Information: K(-) = Kontrol negatif; K(+) = Kontrol positif; P1 = *A. bilimbi* L. concentrations of 25%; P2 = *A. bilimbi* L. concentrations of 50%; P3 = *A. bilimbi* L. concentrations of 75%; and P4 = *A. bilimbi* L. concentrations of 100%. In the figure there is an inhibition carried out by *A. bilimbi* L extract. against *S. typhi*. A measure of inhibitory power is used to determine the ability of *A. bilimbi* L extract.

DISCUSSION

Averrhoa bilimbi L. (Oxalidaceae) is a commonly cultivated and planted plant species in Indonesia. The fruit of *A. bilimbi* L. has a highly sour flavor due to the high concentration of oxalic acid, which exceeds 70% ¹¹. Additionally, the fruit of *A. bilimbi* L. includes bioactive substances that may be utilized to make herbal medication (*Usada*), including saponins, flavonoids, triterpenoids, tannins, and peroxides, formic acid, glucose, calcium oxalate, and sulfur ¹². The fruit's bioactive chemicals were shown to inhibit the growth of disease-causing bacteria, including *Salmonella typhi*, which causes typhoid fever.

According to the results of testing the antibacterial activity of *A. bilimbi* L. fruit extract at a 25% concentration, it failed to inhibit the growth or destruction of *S. typhi* bacteria. This occurred because the concentration of bioactive compounds in the extract was insufficient to inhibit the growth or destruction of the bacterial cell wall, rendering it ineffective at this concentration. This is consistent with previous studies, which demonstrated that bioactive chemicals could suppress infections if they are present in appropriate amounts and proportions and are capable of causing harm to target organs through specific pathways ¹³.

Furthermore, at a concentration of 50%, an average inhibition of $13,000 \pm 1,414$ mm was obtained, and a 75% concentration of an average inhibition of $18,750 \pm 1,500$ with decisive criteria. Meanwhile, at 100% concentration, it inhibited 20.250 ± 1.707 mm with decisive criteria. The high level of inhibition carried out by the extract of *A. bilimbi* L. was due to the presence of bioactive compounds that worked and formed an inhibition zone which became an indicator that the extract with this concentration had been able to inhibit the bacteria available on MHA media ^{11,14,15}. The results showed that the inhibitory capacity of the extract of *A. bilimbi* L. depended on the concentration used. The greater the extract concentration given, starting from a concentration of 50%, 75%, and 100%, the greater the inhibition zone formed by *S.typhi*. It indicates that the bioactive compounds in the extract's concentration work well against pathogens ¹⁶⁻¹⁸.

A. bilimbi L. fruit extract is documented to include a variety of bioactive chemicals with antibacterial, antioxidant, antimicrobial, antiaging, and antioxidative effects. ^{19,20}. Furthermore, the antibacterial benefits of *A. bilimbi* L. fruit were obtained because it contains alkaloids, saponins, flavonoids, and tannins ¹⁴. Flavonoid compounds have a toxic effect on bacteria by damaging the bacterial cell wall so that it is

hydrolyzed and unable to sustain life ²¹. Alcohol groups in flavonoids will react with lipid compounds and amino acids in bacteria so that the bacterial cell wall will be damaged ^{22,23}. Tannin compounds in the fruit extract of *A. bilimbi* L. work by lysing cell walls, inactivating enzymes, and genetic functions of *S. thypi* bacteria ⁹. This is in line with previous research that stated that the ethyl acetate compound in *A. bilimbi* L. fruit which was formulated in the form of gel and extract, was able to provide an excellent inhibitory effect on *Propionibacterium acnes* and *S. aureus* so that it was effectively used as an antibacterial and antimicrobial agent ^{17,24}. The essential oil contained and produced from the fruit extract of *A. bilimbi* L. is helpful as an antibacterial, antioxidant, and *antibiofilm* ¹⁶.

Identifying flavonoid and *triterpenoid* compounds in plant parts is generally done by dissolving certain substances ²⁵. The orange color is formed to detect flavonoid compounds and reddish-purple to detect *triterpenoid* compounds ^{18,26}. Flavonoids in sufficient concentrations can inhibit the growth of *S. thypi* by forming complex compounds with extracellular and dissolved proteins, causing damage to the bacterial cell membrane, followed by the release of effective intracellular compounds that are effective and have antibacterial potential ¹⁹. Meanwhile, the *triterpenoid* content reacts with the *porin* (transmembrane protein) on the

outer membrane of the bacterial cell wall, forming a polymeric solid bond that results in bacteria lysis and growth arrest; this is why the triterpenoid content in the fruit of *A. bilimbi* L. is used as an antiradical or antioxidant^{27,28}.

In vitro and molecular docking studies have been conducted to determine the effectiveness of chemicals found in *A. bilimbi* L. fruit, particularly polyphenol compounds often employed as antibacterial and antioxidant agents in daily life²⁹. Triterpene saponin chemicals found in *A. bilimbi* L. fruit operate as antimicrobials by affecting the integrity of bacterial cell membranes, which results in cell wall lysis.²³ Saponins are semipolar compounds that are soluble in lipids and water, which can lead to the concentration of substances in microbial cells^{11,30}. Concentrating the extract to a sufficiently high concentration enabled the fruit extract of *A. bilimbi* L. to be beneficial as an antibacterial, particularly for its inhibitory efficacy against *S. typhi*. A concentration of 100% can block the bacterium *S. typhi*, which meets the rigorous requirements for *Ciprofloxacin*.

CONCLUSION

At doses of 50%, 75%, and 100%, the fruit extract of *Averrhoa bilimbi* L. (Oxalidaceae) effectively inhibits the bacteria *Salmonella typhi*, which causes *typhoid* fever. The highest extract concentration has the most substantial inhibitory effect and is equal to Ciprofloxacin (decisive criteria). Further study is required to determine the composition, chemical profile, and bioactive chemicals employed as antibacterial and antimicrobial agents, particularly in reducing the development of *Salmonella typhi* bacteria. Experimental animal testing is strongly recommended to establish a more thorough effect.

ACKNOWLEDGEMENT

The authors like to express their gratitude to the Head of Department/KSM Clinical Microbiology, Udayana University, for providing and facilitating the completion of this study. Furthermore, to everyone who assisted and contributed to the seamless operation of this research.

REFERENCES

1. Purba IE, Wandra T, Nugrahini N, Nawawi S, Kandun N. Typhoid

- Fever Control Program in Indonesia: Challenges and Opportunities. *Media Penelit dan Pengemb Kesehat.* 2016;26(2):99–108.
2. Adnyana IMDM, Sudaryati NLG, Suardana A. K. Blood Smear Profile of Patients With Dengue Hemorrhagic Fever in Bali Royal Hospital. *J Vocat Heal Stud.* 2021;5(1):39.
 3. Ministry of Health. *Indonesia Health Profile 2019.* Jakarta: Ministry of Health Republic of Indonesia; 2020. p. 487.
 4. Kumar A, Kumar A. Antibiotic resistome of *Salmonella typhi*: molecular determinants for the emergence of drug resistance. *Front Med.* 2021;15(5):693–703.
 5. Manesh A, Meltzer E, Jin C, Britto C, Deodhar D, Radha S, et al. Typhoid and paratyphoid fever: A clinical seminar. *J Travel Med.* 2021;28(3):1–13.
 6. Prastiyanto ME, Wardoyo FA, Wilson W, Darmawati S. Antibacterial Activity of Various Extracts of *Averrhoa bilimbi* against Multidrug-Resistant Bacteria. *Biosaintifika J Biol Biol Educ.* 2020;12(2):163–8.
 7. Kimbal AVR, Jatmiko YD, Ardyati T. Antimicrobial Activity of Combination Bacteriocin and Asam Suntik Extract (*Averrhoa bilimbi* L. fermented) Against Multidrug-Resistant *Escherichia coli* in Lettuces (*Lactuca sativa*). *J Exp Life Sci.* 2021;11(2):60–7.
 8. Iwansyah AC, Desnilasari D, Agustina W, Pramesti D, Indriati A, Mayasti NKI, et al. Evaluation on the physicochemical properties and mineral contents of *Averrhoa bilimbi* L. Leaves dried extract and its antioxidant and antibacterial capacities. *Food Sci Technol.* 2021;41(4):987–92.
 9. Khoo HE, Azlan A, Kong KW, Ismail A. Phytochemicals and Medicinal Properties of Indigenous Tropical Fruits with Potential for Commercial Development. *Evidence-Based Complement Altern Med.* 2016;2016(7591951):1–20.
 10. Darwin M, Mamondol MR, Sormin SA, Nurhayati Y, Tambunan H, Sylvia D, et al. *Metode penelitian pendekatan kuantitatif (Quantitative approach research method).* 1st ed. Bandung: CV Media Sains Indonesia; 2021. 192 p.
 11. Fidrianny I, Rahmawati A, Hartati R. Comparison profile of different extracts of *Averrhoa bilimbi* L. In antioxidant properties and phytochemical content. *Rasayan J Chem.* 2018;11(4):1628–34.
 12. Prasathkumar M, Anisha S, Dhriya C, Becky R, Sadhasivam S. Therapeutic and pharmacological efficacy of selective Indian

- medicinal plants – A review. *Phytomedicine Plus*. 2021;1:100029.
13. Anuar NA, Salleh RM. Development of fruit jam from *Averrhoa bilimbi* L. *J Food Process Preserv*. 2019;43(e13904):1–7.
 14. Xoca-Orozco LA, Zamora-Gasga V, Espinosa-Alonso G, Velázquez-Estrada RM, López-García U, Sáyago-Ayerdi S, et al. In vitro antioxidant and antifungal activities of carambola (*Averrhoa carambola* L), extracts. *Biocatal*. 2018;20(2):104–9.
 15. Raja Arifin RNA, Jumal J. Antioxidant Activity of *Averrhoa bilimbi* Linn. Leaves Extract Using Two Different Types of Solvents. *Malaysian J Sci Heal Technol*. 2021;7:76–82.
 16. Nartey D, Accorley ED, Opoku R, Darko G, Borquaye LS. Essential Oils from *Averrhoa carambola* L. (Oxalidaceae): Chemical Composition, Antioxidant, Antimicrobial, and Anti-biofilm Potential. *Chem Africa*. 2021;4(4):741–52.
 17. Simanjuntak HA, Gurning K, Sinaga VB. Antibacterial Activity of Cold Powder Preparation of (Ethanol Extract) Starfruit Leaf (*Averrhoa bilimbi* Linn.) Against *Propionibacterium acnes*. *J Pembelajaran Dan Biol Nukl*. 2020;6(2):120–8.
 18. Leliqia NPE, Safitri I. A review of phytochemical properties, antibacterial activity, and toxicity study of *Averrhoa bilimbi* leaves and fruit. *J Pharm Sci Appl*. 2021;3(1):32–9.
 19. Seebaluck-Sandoram R, Lall N, Fibrich B, Blom van Staden A, Saleem H, Mahomoodally MF. Antimicrobial, antioxidant, and cytotoxic evaluation of two underutilized food plants: *Averrhoa bilimbi* L. (Oxalidaceae) and *Phyllanthus acidus* L. Skeels (Phyllanthaceae). *Biocatal Agric Biotechnol*. 2019;18(December 2018):100998.
 20. Hasim H, Arifin YY, Andrianto D, Faridah DN. Ethanol Extracts of *Averrhoa Bilimbi* Leaf Demonstrated Antioxidative and Anti-inflammatory Activity. *J Apl Teknol Pangan*. 2019;8(3):86–93.
 21. Nakhil U, Sikumbang IM, Hendra Putri N, Lutfiyati H. Wuluh Starfruit (*Averrhoa Bilimbi*) Extract Gel For Recurrent Aftosa Stomatitis. *J Farm Sains dan Prakt*. 2019;5(2):2579–4558.
 22. Samputri RD, Toemon AN, Widayati R. Antibacterial activity test of ethanol extract of kamandrah seeds (*Croton tiglium* L.) Against the Growth of *Salmonella typhi* with Disc Diffusion Method (Kirby-Bauer). *Herb-Medicine J*. 2020;3(3):19–33.
 23. Abuga I, Sulaiman SF, Abdul Wahab R, Ooi KL, Abdull Rasad MSB. Phytochemical constituents and antibacterial activities of 45

- Malay traditional medicinal plants. *J Herb Med.* 2022;32(July 2021):100496.
24. Pertiwi D, Hafiz I, Jannah W, Winata HS, Sari M, Suroyo RB. Antibacterial activities of belimbing wuluh (*Averrhoa bilimbi* L.) ethyl acetate extract on gel formulated against *propionibacterium acnes* and *staphylococcus aureus*. *Int J Appl Pharm.* 2020;12(6):224–8.
 25. Averrhoa F, Rante H, Yasir Y, Nur S, Emba A. Detection of Antimicrobial Compounds by Bioautography from Star Fruit (*Averrhoa bilimbi* Linn.). *J Pharm Med Sci.* 2018;3(1):1–5.
 26. Setyawan HY, Sukardi S, Nareswari BF. The phytochemical potential of *Averrhoa bilimbi* - A review. *IOP Conf Ser Earth Environ Sci.* 2021;733:012091.
 27. Ahmed Q, Alhassan A, Khatib A, Shah S, Hasan M, Sarian M. Antiradical and Xanthine Oxidase Inhibitory Activity Evaluations of *Averrhoa bilimbi* L. Leaves and Tentative Identification of Bioactive Constituents through LC-QTOF-MS/MS and Molecular Docking Approach. *Antioxidants.* 2018 Oct 8;7(10):137.
 28. Adnyana IMDM, Sudaryati NLG, Sitepu I. Toxicity of Legiayu incense as Insecticide and Larvicide Against *Aedes aegypti* Mosquitoes Mortality. *Indones J Pharm.* 2021;32(4):524–521.
 29. Ullah H, Hussain Y, Santarcangelo C, Baldi A, Di Minno A, Khan H, et al. Natural Polyphenols for the Preservation of Meat and Dairy Products. *Molecules.* 2022;27(6):1906.
 30. Pang D, Liao S, Wang W, Mu L, Li E, Shen W, et al. Destruction of the cell membrane and inhibition of cell phosphatidic acid biosynthesis in: *Staphylococcus aureus*: An explanation for the antibacterial mechanism of morusin. *Food Funct.* 2019;10(10):6438–46.